

Laboratório de Hantaviroses e Rickettsioses

2006

Indexado (fator de impacto > 0,65 a 2,0)

Rozental T, Eremeeva ME, Paddock CD, Zaki SR, Dasch GA, Lemos ERS 2006. Fatal case of Brazilian spotted fever confirmed by immunohistochemical staining and sequencing methods on fixed tissues.. Ann NY Acad Sci 1078: 257-259.

2007

Indexado (fator de impacto <0,65 e não impactadas no ISI)

Lamas CC, Favacho ARM, Santos M, Rozental T, Boia MN, Lemos ERS 2007. Bartonella native valve endocarditis: the first Brazilian case alive and well. Braz J Infect Dis 11: 591-594.

Albuquerque JI, Gonçalves AQ, Lemos ERS 2007. Intestinal protozoans and helminths among Terena Indians in the State of Mato Grosso do Sul, Brazil: High prevalence of Blastocystis hominis. Rev Soc Bras Med Trop 40: 631-634.

Luz CF, Thomé ML, Barth OM 2007. Recursos tróficos de Aipis mellifera (Hymenoptera, Apidae) na região de Morro Azul do Tingúá, estado do Rio de Janeiro. Braz J Bot 30: 29-36.

Indexado (fator de impacto > 0,65 a 2,0)

Barreto DF, Takiya CM, Schatzmayr HG, Nogueira RMR, Filho JCF, Barth OM 2007. Histopathological and ultrastructural aspects of mice lungs experimentally infected with dengue virus serotype 2. Mem I Oswaldo Cruz (impresso) 102: 175-182.

Barreto CF, Luz CFP, Baptista-neto JA, Vilela CG, Barth OM 2007. Palynological analysis of a sediment core obtained in Guanabara Bay, Rio de Janeiro, Brazil. An Acad Bras Cienc 79: 223-234.

Indexado (fator de impacto >2,0 a 4,0)

Raboni SM, Levis S, Rosa ES, Bisordi I, Delfraro A, Lemos E, Correia DC, Duarte dos Santos CN 2007. Hantavirus infection in Brazil: development and evaluation of an enzyme immunoassay and immunoblotting based on N recombinant protein. Diagn Micr Infec Dis 58: 89-97.

2008

Indexado (> 0,65 a 2,0)

doi:> Lemos ERS, Lamas CC, Curi A 2008. Human bartonellosis: seroepidemiological and clinical aspects with an emphasis on data from Brazil - a review. Mem I Oswaldo Cruz (impresso) 103: 221-235.

doi:> Coelho LG, Barth OM, Araújo DS 2008. Pollen analysis of holocene sediments from the Poço das Antas National Biological Reserve, Silva Jardim, Rio de Janeiro, Brazil. An Acad Bras Cienc 80: 531-541.

Indexado (>4,0)

doi:> Martins AN, Medeiros SO, Simonetti JP, Schatzmayr HG, Tanuri A, Brindeiro RM 2008. Phylogenetic and genetic analysis of feline immunodeficiency virus gag, pol, and env genes from domestic cats undergoing

nucleoside reverse transcriptase inhibitor treatment or treatment-naïve cats in Rio de Janeiro, Brazil. J Virol 82: 7863-7874.

Indexado (<0,65 e não impactadas no ISI)

doi: Pinto Jr VL, Curi A, Rozental T, Lemos ERS 2008. Cat scratch disease complicated with aseptic meningitis and neuroretinitis. Braz J Infect Dis 12: 158-160.

doi: Lemos ER, Lamas CC, Favacho AR, Rozental T, Bóia MN, Silva AG, Barreiras J 2008. Characterization of Rickettsia rickettsii in a case of fatal Brazilian spotted fever in the city of Rio de Janeiro, Brazil. Braz J Infect Dis 12: 149-151.

doi: Munhoz M, Luz CF, Messner Filho PE, Barth OM, Reinert F 2008. Viabilidade polínica de Carica papaya L.: uma comparação metodológica. Braz J Bot 31: 209-214.

Indexado (>2,0 a 4,0)

doi: Portz C, Beltrão N, Furian TQ, Biano-júnior A, Macagnan M, Griebeler J, Rosa CAVL, Colodel EM, Driemeier D, Back A, Barth OM, Canal CW 2008. Natural infection of turkeys by infectious laryngotracheitis virus. Vet Microbiol 131: 57-64.

2009

Indexado (fator de impacto > 0,65 e < 2)

doi: Oliveira RC, Teixeira BR, Mello FC, Pereira AP, Duarte AS, Bonaldo MC, Bonvicino CR, d'Andrea PS, Lemos ER 2009. Genetic characterization of a Juquitiba-like viral lineage in Oligoryzomys nigripes in Rio de Janeiro, Brazil. Acta Trop 112: 212-218.

Indexado (fator de impacto > 2 e < 4)

doi: Raboni SM, Hoffmann FG, Oliveira RC, Teixeira BR, Bonvicino CR, Stella V, Carstensen S, Bordignon J, d'Andrea PS, Lemos ER, dos Santos CN 2009. Phylogenetic characterization of hantaviruses from wild rodents and hantavirus pulmonary syndrome cases in the state of Paraná (southern Brazil). J Gen Virol 90: 2166-2171.

doi: Rozental T, Favacho AR, Barreira JD, Oliveira RC, Gomes R, Almeida DN, Lemos ER 2009. Rickettsia spp. infection in *Rhipicephalus sanguineus* ticks in a Brazilian spotted fever endemic rural area in Rio de Janeiro state, Brazil. Clin Microbiol Infec 15: 245-246.

doi: Lamas CC, Rozental T, Boia MN, Favacho AR, Kirsten AH, da Silva AP, de Lemos ER 2009. Seroprevalence of *Coxiella burnetii* antibodies in human immunodeficiency virus-positive patients in Jacarepaguá, Rio de Janeiro, Brazil. Clin Microbiol Infec 15 suppl 2: 140-141.

Indexado (fator de impacto > 4)

doi: Limongi JE, da Costa FC, Pinto RM, de Oliveira RC, Bragagnolo C, Lemos ER, de Paula MB, Pajuaba Neto AA, Ferreira MS 2009. Cross-sectional survey of hantavirus infection, Brazil. Emerg Infect Dis 15: 1981-1983.

2010

Indexado (fator de impacto < 0,65 ou sem FI)

doi: Curi AL, Machado D, Heringer G, Campos WR, Lamas C, Rozental T, Gutierrez A, Orefice F, Lemos E 2010. Cat-scratch disease: ocular manifestations and visual outcome. *Int Ophthalmol* 30: 553-558.

Indexado (fator de impacto > 0,65 e < 2)

doi: Lemos ER, Mares-Guia MA, de Almeida DN, da Silva RG, Silva CM, Britto C, Lamas CC 2010. Febre do viajante associada com adenite cervical e sororreatividade para Bartonella sp em paciente brasileira, após retorno da África do Sul. *Rev Soc Bras Med Trop* 43: 472-473.

Indexado (fator de impacto > 2 e < 4)

doi: Lamas CC, Mares-Guia MA, Rozental T, Moreira N, Favacho AR, Barreira J, Guterres A, Boia MN, de Lemos ER 2010. Bartonella spp. infection in HIV positive individuals, their pets and ectoparasites in Rio de Janeiro, Brazil: serological and molecular study. *Acta Trop* 115: 137-141.

Indexado (fator de impacto > 4)

doi: Cristiane Lamas 2010. Diagnostic strategy for blood culture-negative endocarditis. *Clin Infect Dis* 51: 141-142.

doi: Travassos da Rosa ES, Lemos ER, Medeiros DB, Simith DB, Pereira AS, Elkhoury MR, Mendes WS, Vidigal JR, de Oliveira RC, D'Andrea PS, Bonvicino CR, Cruz AC, Nunes MR, Pedro Vasconcelos 2010. Hantaviruses and hantavirus pulmonary syndrome, Maranhão, Brazil. *Emerg Infect Dis* 16: 1952-1955.

2011

Indexado (fator de impacto > 2 e 4)

doi: de Oliveira RC, Padula PJ, Gomes R, Martinez VP, Bellomo C, Bonvicino CR, Freire e Lima DI, Bragagnolo C, Caldas AC, D'Andrea PS, Lemos ER 2011. Genetic characterization of hantaviruses associated with sigmodontine rodents in an endemic area for hantavirus pulmonary syndrome in southern Brazil. *Vector-Borne Zoonot* 11: 301-314.

doi: Lemos ER, Rozental T, Mares-Guia MA, Almeida DN, Moreira N, Silva RG, Barreira JD, Lamas CC, Favacho AR, Damasco PV 2011. Q fever as a cause of fever of unknown origin and thrombocytosis: first molecular evidence of Coxiella burnetii in Brazil. *Vector-Borne Zoonot* 11: 85-87.

Indexado (fator de impacto > 4)

doi: Silva N, Eremeeva ME, Rozental T, Ribeiro GS, Paddock CD, Ramos EA, Favacho AR, Reis MG, Dasch GA, de Lemos ER, Ko AI 2011. Eschar-associated spotted fever rickettsiosis, Bahia, Brazil. *Emerg Infect Dis* 17: 275-278.

doi: Travassos da Rosa ES, Medeiros DB, Nunes MR, Simith DB, de Souza Pereira A, Elkhoury MR, Lavocat M, Marques AA, Via AV, D'Andrea P, Bonvicino CR, Lemos ER, Vasconcelos PF 2011. Pygmy rice rat as potential host of Castelo dos Sonhos hantavirus. *Emerg Infect Dis* 17: 1527-1530.

Indexado (fator de impacto 0,65 e 2)

doi: Damasco PV, Ávila CA, Barbosa AT, Carvalho MM, Pereira GM, Lemos ER, Boia MN, Pereira MM 2011. Atypical lymphocytosis in leptospirosis: a cohort of hospitalized cases between 1996 and 2009 in State of Rio de Janeiro, Brazil. *Rev Soc Bras Med Trop* 44: 611-615.

doi: Favacho AR, Rozental T, Calic SB, Scofield MA, Lemos ER 2011. Fatal Brazilian spotless fever caused by Rickettsia rickettsii in a dark-skinned patient. *Rev Soc Bras Med Trop* 44: 395-396.

doi: Crissiuma A, Favacho A, Gershony L, Mendes-de Almeida F, Gomes R, Mares-Guia MA, Rozental T, Barreira J, Lemos E, Labarthe N 2011. Prevalence of Bartonella species DNA and antibodies in cats (*Felis catus*) submitted to a spay/neuter program in Rio de Janeiro, Brazil. *J Feline Med Surg* 13: 149-151.

2012

Indexado (fator de impacto >=1 e < 2,71)

doi: Rozental T, Mascarenha LF, Rozenbaum R, Gomes R, Mattos GS, Magno CC, Almeida DN, Rossi MI, Favacho AR, de Lemos ER 2012. *Coxiella burnetii*, the agent of Q fever in Brazil: its hidden role in seronegative arthritis and the importance of molecular diagnosis based on the repetitive element IS1111 associated with the transposase gene. *Mem I Oswaldo Cruz (impresso)* 107: 695-697.

doi: de Oliveira RC, Guterres A, Schrago CG, Fernandes J, Teixeira BR, Zeccher S, Bonvicino CR, D'Andrea PS, de Lemos ER 2012. Detection of the first incidence of Akodon paranaensis naturally infected with the Jabora virus strain (Hantavirus) in Brazil. *Mem I Oswaldo Cruz (impresso)* 107: 424-428.

doi: Raboni SM, Delfraro A, de Borba L, Teixeira BR, Stella V, de Araujo MR, Carstensen S, Rubio G, Maron A, Lemos ER, D'Andrea PS, Duarte dos Santos CN 2012. Hantavirus infection prevalence in wild rodents and human anti-hantavirus serological profiles from different geographic areas of South Brazil. *Am J Trop Med Hyg* 87: 371-378.

Indexado (fator de impacto >=4,35 e < 7,51)

doi: Travassos da Rosa ES, Medeiros DB, Nunes MR, Simith DB, Pereira AS, Elkhoury MR, Santos ED, Lavocat M, Marques AA, Via AV, Kohl VA, Terças AC, D'Andrea P, Bonvicino CR, Sampaio de Lemos ER, Vasconcelos PF 2012. Molecular epidemiology of Laguna Negra virus, Mato Grosso State, Brazil. *Emerg Infect Dis* 18: 982-985.

2013

Indexado (fator de impacto < 1 ou sem FI)

doi: Marcus Pires , Santos TM, Santos HA, Vilela JA, Roier EC, Peixoto MP, Silva CB, Jairo Dias Barreira, Lemos ERS, Carlos Massard 2013. Amblyomma cajennense infestation on horses in two microregions of the state of Rio de Janeiro, Brazil.. *Rev Bras Med Vet* 22: 235-242.

Indexado (fator de impacto >=1 e < 2,71)

cristiane lamas, Santos RCOP, Silva RG, Vicente-de-Mattos LHB, Barros-de-Almeida E, Lemos ERS, Boia MN 2013. Hantavirus infection in HIV positive individuals in Rio de Janeiro, Brazil: a seroprevalence study. *Braz J Infect Dis* 17: 120-121.

Indexado (fator de impacto >=2,71 e < 4,35)

doi: Santos RCOP, Gentile R, Silva AG, Fernandes-de-Jesus J, Teixeira BR, Vaz VV, Valdez FP, Vicente-de-Mattos LHB, Costa-Neto SF, Bonvicino CR, D'Andrea PS, Lemos ERS 2013. Ecological study of hantavirus infection in wild rodents in an endemic area in Brazil.. *Acta Trop* 131C: 1-10.

doi: Silva AG, Santos RCOP, Fernandes-de-Jesus J, D'Andrea PS, Bonvicino CR, Bragnolo C, Guimarães GD, Almada GL, Machado PR, Lavocat M, Elkhoury M, Schrago C, Lemos ERS 2013. Phylogenetic analysis of the S segment from Juquitiba hantavirus: Identification of two distinct lineages in *Oligoryzomys nigripes*. *Infect Genet Evol* 18: 262-268.

2014

Indexado (fator de impacto < 1 ou sem FI)

doi: Silva AG, Santos RCOP, Fernandes-de-Jesus J, Pereira LS, Silva FCD, Oliveira FCG, D'Andrea PS, Bonvicino CR, Carlos Guerra Schrago, Lemos ERS 2014. Characterization of Juquitiba Virus in Oligoryzomys fornesi from Brazilian Cerrado. *Viruses* 6: 1473-1482.

doi: Santos RCOP, Silva AG, Fernandes-de-Jesus J, D'Andrea PS, Bonvicino CR, Lemos ERS 2014. Hantavirus Reservoirs: Current Status with an Emphasis on Data from Brazil. *Viruses* 6: 1929-1973.

doi: Favacho ARM, Roger I, Amanda Kitada, Pessoa Junior AA, Silva RG, Tupy-de-Godoy D, Sandro Pereira, Lemos ERS 2014. MOLECULAR IDENTIFICATION OF Bartonella henselae IN A SERONEGATIVE CAT SCRATCH DISEASE PATIENT WITH AIDS IN RIO DE JANEIRO, BRAZIL. *Rev Inst Med Trop SP* 56: 363-365.

Mares-guia MAMM, Rozental T, Silva AG, Silva RG, Peixoto-de-Almeida DN, Moreira NS, Barreira JD, Favacho ARM, Andrea Lopes Santana, Lemos ERS 2014. Molecular identification of the agent of Q fever – Coxiella burnetii – in domestic animals in State of Rio de Janeiro, Brazil.. *Rev Soc Bras Med Trop* 47: 231-234.

Cunha, N.C., Lemos ERS, Rozental T, Teixeira R.C., Cordeiro, M.D., Lisboa, R.S., Favacho ARM, Barreira JD, Rezende J, Fonseca A.H. 2014. Rickettsiae of the Spotted Fever Group in dogs, horses and ticks: an epidemiological study in an endemic region of the state of Rio de Janeiro, Brazil.. *Rev Bras Med Vet* 36: 294-300.

Indexado (fator de impacto >=1 e < 2,71)

doi: Rozental T, Ferreira MS, Silva RG, Carolina Monteiro, Paulo Barbosa, Itacirema Bezerra, Márcio Garcia, Rafael Galliez, Stefan Oliveira, Patrícia Brasil, Tatiana Resende, Lemos ERS 2014. A cluster of Rickettsia rickettsii infection at an animal shelter in an urban area of Brazil.. *Epidemiol Infect* 8: 1-5.

doi: Santos RCOP, Gentile R, Silva AG, Fernandes-de-Jesus J, Teixeira BR, Vanderson Vaz, Fernanda Pedone Valdez, Vicente-de-Mattos LHB, Costa-Neto SF, Bonvicino CR, D'Andrea PS, Lemos ERS 2014. Ecological study of hantavirus infection in wild rodents in an endemic area in Brazil. *Acta Trop* 131: 1-10.

doi: Vitor Laerte Pinto Jr, A Souza, Lemos ERS 2014. Regional variations and time trends of hantavirus pulmonary syndrome in Brazil.. *Epidemiol Infect* 7: 1-6.

Indexado (fator de impacto >=2,71 e < 4,35)

doi: Teixeira BR, Nathalie Loureiro, Pereira LS, Gentile R, Santos RCOP, Silva AG, Fernandes-de-Jesus J, Vicente-de-Mattos LHB, Sônia Raboni, Giselia Rubio, Cláudia Duarte, Bonvicino CR, Lemos ERS, D'Andrea PS 2014. Population Ecology of Hantavirus Rodent Hosts in Southern Brazil. *Am J Trop Med Hyg* 13: 0465-0471.

doi: Lívia de Barros Lopes, Silva AG, Rozental T, Santos RCOP, Mares-guia MAMM, Fernandes-de-Jesus J, José Ferreira Figueiredo, Inês Anshau, Sebastião de Jesus, Ana Beatriz Almeida, Valéria Cristina da Silva, Alba Valéria Gomes de Melo Via, Bonvicino CR, D'Andrea PS, Barreira JD, Lemos ERS 2014. Rickettsia bellii, Rickettsia amblyommii, and Laguna Negra hantavirus in an Indian reserve in the Brazilian Amazon. *Parasite Vector* 7: - .

Indexado (fator de impacto >=4,35 e < 7,51)

doi: Santos RCOP, Marcelo Cordeiro Santos, Silva AG, Fernandes-de-Jesus J, X.Mello, AP João G, Bernardino Claudio Albuquerque, Pedro Vasconcelos, Lemos ERS 2014. Rio Mamore Virus and Hantavirus Pulmonary

Syndrome, Brazil.. *Emerg Infect Dis* 20: 1568-1570.

2015

Indexado (fator de impacto >=1 e < 2,71)

doi: Rozental T, Ferreira MS, Silva RG, Carolina Monteiro, Paul Roberto Barbosa, Itacirema Bezerra, Rafael Galliez, Patricia Brasil, Lemos ERS 2015. A cluster of *Rickettsia rickettsii* infection at an animal shelter in an urban area of Brazil. *Epidemiol Infect* 143: 2446-2450.

doi: Silva AG, Santos RCOP, Fernandes-de-Jesus J, Carlos Guerra Schrago, Lemos ERS 2015. Detection of different South American Hantavirus. *Virus Res* 210: 106-113.

doi: Figueiredo AS, Lampe E, Espírito-Santo MP, Mello FCA, Almeida FQ, Lemos ERS, Godoi TL, Dimache LA, Santos DR, Villar LM 2015. Identification of two phylogenetic lineages of equine hepatic virus and high prevalence in Brazil. *Vet J* 206: 414-416.

doi: Solange Maria Gennari, Ogrzewalska MH, Herbert Sousa Soares, Danilo Gonçalves Saraiva, Adriano Pinter, Fernanda Aparecida Nieri-Bastos, Marcelo B. Labruna, Matias Pablo Juan Azabó, Jitender Prakash Dubey 2015. Toxoplasma gondii antibodies in wild rodents and marsupials from the Atlantic Forest, state of São Paulo, Brazil. *Vet Parasitol* 24: 379-382.

Indexado (fator de impacto >=2,71 e < 4,35)

doi: Ogrzewalska MH, Ivan Literak, Miroslav Capek, Oldrich Sychra, Víctor Álvarez Calderón, Bernardo Calvo Rodríguez, Carlos Prudencio, Thiago F. Martins, Marcelo B. Labruna 2015. Bacteria of the genus *Rickettsia* in ticks (Acaris: Ixodidae) collected from birds in Costa Rica. *Ticks Tick-Borne Dis* 6(4): 478-482.

doi: Fernandes-de-Jesus J, Santos RCOP, Silva AG, Fabiana de Carvalho Serra, Bonvicino CR, D'Andrea PS, Rivaldo Venâncio Cunha, Silvana Levis, Lemos ERS 2015. Co-circulation of Clade C New World Arenaviruses: New geographic distribution and host species. *Infect Genet Evol* 33: 242-245.

doi: Camile Lugarini, Thiago F. Martins, Ogrzewalska MH, Nathália Costa Teixeira de Vasconcelos, Vincenzo A. Ellis, Jacqueline Bianque de Oliveira, Adriano Pinter, Marcelo B. Labruna, Jean Carlos Ramos Silva 2015. Rickettsial agents in avian ixodid ticks in northeast Brazil. *Ticks Tick-Borne Dis* 6: 364-375.

doi: Marketa Novakova, Ivan Literak, Luis Chevez, Thiago F. Martins, Ogrzewalska MH, Marcelo B. Labruna 2015. Rickettsial infections in ticks from reptiles, birds and humans in Honduras. *Ticks Tick-Borne Dis* 6: 737-742.

doi: Ivan Literak, Ana Claudia Norte, Maria Sofia núncio, Isabel Lopes de Carvalho, Ogrzewalska MH, Markéta Nováková, Thiago F. Martins, Oldrich Sychra, Roberto Resendes, Pedro Rodrigues 2015. Ticks on passerines from the Archipelago of the Azores as hosts of borreliae and rickettsiae. *Ticks Tick-Borne Dis* 6: 607-610.

doi: Favacho ARM, Santos RCOP, Bonvicino CR, D'Andrea PS, Lemos ERS 2015. Zoonotic Bartonella species in wild rodents in the state of Mato Grosso do Sul, Brazil.. *Microbes Infect* 17: 889-892.

Qualis A1

doi: Silva AG, Fernandes-de-Jesus J, Santos RCOP, Lemos ERS 2015. Is the evolution of Hantavirus driven by its hist?. *Genetics and Evolution* 35: 142-143.

2016

Indexado (fator de impacto < 1 ou sem FI)

doi: Lupi O, Carvalho E, Rozental T, Favacho ARM, Lemos ERS, Brasil P 2016. Serological Evidence of Human Coinfection by Brazilian Spotted Fever and Bartonellosis. *Int J Clin Med* 7: 766-770.

doi: Ogrzewalska MH, Adriano Pinter 2016. Ticks (Acari: Ixodidae) as ectoparasites of Brazilian Wild birds and their association with rickettsial diseases. *Braz J Vet Res Anim Sci* 53: 1-31.

Indexado (fator de impacto >=1 e < 2,71)

doi: Ogrzewalska MH, Fernanda A. Nieri-Bastos, Arlei Marcili, Santiago Nava, Daniel González-Acuña, Sebastián Muñoz-Leal, Ignacio Ruiz-Arrondo, José M. Venzal, Atilio Mangold, Marcelo B. Labruna 2016. A novel spotted fever group Rickettsia infecting Amblyomma parvitarsum (Acari: Ixodidae) in Highlands of Argentina and Chile. *Ticks Tick-Borne Dis* 7: 439-442.

doi: Ogrzewalska MH, Schwarcz K, Bajay MM, Bajay SK, Pinheiro JB, Zucchi MI, Pinter A, Labruna MB 2016. Characterization of Genetic Variability and Population Structure of the Tick (Acari: Ixodidae). *J Med Entomol* 53: 843-850.

doi: Hermes Luz, McIntosh D, Furusawa GP, Flausino W, Rozental T, Lemos ERS, Landulfo GA, Faccini JL 2016. Infection of Amblyomma ovale with Rickettsia species Atlantic rainforest in Serra do Mar, São Paulo State, Brazil. *Ticks Tick-Borne Dis* 7: 1265-1267.

doi: Blanco CM, Teixeira BR, Silva AG, Santos RCOP, Pereira LS, Ogrzewalska MH, Lemos ERS 2016. Microorganisms in ticks (Acari: Ixodidae) collected on marsupials and rodents from Santa Catarina, Paraná and Mato Grosso do Sul states, Brazil. *Ticks Tick-Borne Dis* 8: 90-98.

doi: Mares-guia MAMM, Rozental T, Silva AG, Michelle Ferreira dos Santos, Renato de Gasperis Botticini, Ana Kely Carolina Terra, Sandro Marraschi, Rosany Bochner, Lemos ERS 2016. Molecular Identifications of Q Fever in Patients with a Suspected Diagnosis of Dengue in Brazil in 2013-2014. *Am J Trop Med Hyg* 94: 1090-1094.

doi: Marcella Gonçalves Coelho, Vanessa do Nascimento Ramos, Jean Ezequiel Limongi, Lemos ERS, Silva AG, Costa-Neto SF, Rozental T, Bonvicino CR, D'Andrea PS, Jonas Moraes-Filho, Marcelo Bahia Labruna, Matias Pablo Juan Szabó 2016. Sorologic evidence of the exposure of small mammals to spotted fever Rickettsia and Rickettsia bellii in Minas Gerais, Brazil. *J Infect Dev Countr* 10: 275-282.

Indexado (fator de impacto >=2,71 e < 4,35)

doi: Gonçalves LR, Favacho ARM, Roque AL, Mendes NS, Fidelis Junior OL, Benevenute JL, Herrera HM, D'Andrea PS, Lemos ERS, Zacaria R, André MR 2016. Association of Bartonella Species with Wild and Synanthropic Rodents in Different Brazilian Biomes. *Appl Environ Microb* 82: 7154-7164.

doi: Epelboin L, Nacher M, Mahamat A, Pommier de Santi V, Berlioz-Arthaud A, Eldin C, Abboud P, Briolant S, Mosnier E, Mendonça Gomes Mdo S, Vreden SG, Pierre-Demar M, Lacerda M, Raoult D, Lemos ERS, Djossou F 2016. Q Fever in French Guiana: Tip of the Iceberg or Epidemiological Exception?. *Plos Neglect Trop D* 10: - .

2017